

Dr. Scholl's® ClearAway® Wart Remover

The Drug labeling information on this website may differ from the product labeling on purchased product. Please consult your purchased package for information specific to that product.

Drug Facts	
Active ingredient Salicylic acid 40%	Purpose Wart remover
Uses ▪ for the removal of common warts. The common wart is easily recognized by the rough 'cauliflower-like' appearance of the surface	
Warnings For external use only	
Do not use <ul style="list-style-type: none"> ▪ if you are a diabetic ▪ if you have poor blood circulation ▪ on irritated skin or any area that is infected or reddened ▪ on moles, birthmarks, warts with hair growing from them, genital warts, or warts on the face or mucous membranes 	
Stop use and ask a doctor if discomfort lasts	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions <ul style="list-style-type: none"> ▪ wash affected area ▪ may soak wart in warm water for 5 minutes ▪ dry area thoroughly ▪ if necessary, cut medicated disc to fit wart ▪ apply medicated disc with sticky side adhering to skin ▪ cover medicated disc with enclosed cushion ▪ repeat procedure every 48 hours as needed (until wart is removed) for up to 12 weeks 	
Other information <ul style="list-style-type: none"> ▪ store between 20° to 25°C (68° to 77°F) ▪ protect from excessive heat (temperatures above 40°C (104°F)) 	
Inactive ingredients antioxidant (CAS 991-84-4), iron oxides, mineral oil, petroleum hydrocarbon resin, silicon dioxide, synthetic polyisoprene rubber, talc	
Questions? 1-866-360-3226	