

## Dr. Scholl's® Ingrown Toenail Pain Reliever

The Drug labeling information on this website may differ from the product labeling on purchased product. Please consult your purchased package for information specific to that product.

<b>Drug Facts</b>
<b>Active ingredients</b> <b>Purpose</b> Sodium sulfide 1%.....Ingrown toenail reliever
<b>Use</b> <ul style="list-style-type: none"> <li>▪ for temporary relief of pain and discomfort from ingrown toenails</li> </ul>
<b>Warnings</b> <b>For external use only</b>
<b>Do not use</b> <ul style="list-style-type: none"> <li>▪ on open sores</li> </ul>
<b>Ask a doctor before use if you have</b> <ul style="list-style-type: none"> <li>▪ diabetes</li> <li>▪ poor blood circulation</li> <li>▪ gout</li> </ul>
<b>When using this product</b> <ul style="list-style-type: none"> <li>▪ use with a retainer ring</li> <li>▪ avoid contact with eyes. If product gets in eyes, flush with water for 15 minutes and get medical help right away.</li> </ul>
<b>Stop use and ask a doctor if</b> <ul style="list-style-type: none"> <li>▪ redness or swelling of your toe increases</li> <li>▪ discharge is present around the nail</li> <li>▪ symptoms last more than 7 days or clear up and occur again within a few days</li> </ul>
<b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.
<b>Directions</b> <ul style="list-style-type: none"> <li>▪ adults and children 12 years and over:           <ul style="list-style-type: none"> <li>○ wash affected area and dry thoroughly</li> <li>○ place retainer ring on toe with slot over the area where the ingrown nail and the skin meet. Smooth ring down firmly.</li> <li>○ cut open tube on score mark. Apply enough gel product to fill the slot in the ring. Immediately replace cap on tube.</li> <li>○ place round center section of bandage directly over the gel-filled ring to seal the gel in place. Smooth ends of bandage strip around toes.</li> <li>○ repeat twice daily (morning and night) for up to 7 days until pain and discomfort are relieved or until the nail can be lifted out of the nail groove and easily trimmed</li> </ul> </li> </ul>



Bayer HealthCare  
Consumer Care

- children under 12 years: ask a doctor

***Other information***

- save carton for full labeling
- keep tube tightly closed when not in use
- store between 20° to 25° C (68° to 77° F)

***Inactive ingredients***

edetate disodium, hydroxyethyl cellulose, potassium acetate, water

***Question?*** 1-866-360-3226